



Correction to: COVID-19 in children: clinical and epidemiological spectrum in the community

César Garcia-Vera¹ · Sandra Castejon-Ramirez² · Elena Laín Miranda³ · Rebeca Hernández Abadía² · María García Ventura⁴ · Elena Borque Navarro⁵ · Patricia Rubio Sánchez² · Álvaro Baeta Ruiz⁵ · José María Mengual Gil⁶

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: European Journal of Pediatrics
<https://doi.org/10.1007/s00431-021-04235-4>

In the original published version of this article, the presented Table 4 has incorrect presentation of data. The following table is now presented correctly.

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s00431-021-04235-4>.

✉ César Garcia-Vera
cgarciavera@gmail.com

Sandra Castejon-Ramirez
scastejonr@gmail.com

Elena Laín Miranda
elenalainm@gmail.com

Rebeca Hernández Abadía
herabare@gmail.com

María García Ventura
mariagarciaventura90@gmail.com

Elena Borque Navarro
elenaborna@gmail.com

Patricia Rubio Sánchez
patrirub92@hotmail.com

Álvaro Baeta Ruiz
a.baeta11@gmail.com

José María Mengual Gil
pepebobes@gmail.com

- 1 Primary Health Care Center “José Ramón Muñoz Fernández”, Zaragoza, Spain
- 2 Hospital Materno-Infantil Miguel Servet, Zaragoza, Spain
- 3 Public Health Technician in the Health Region II of Zaragoza, Zaragoza, Spain
- 4 Primary Health Care Center “Valdespartera”, Zaragoza, Spain
- 5 Hospital Clínico Universitario Lozano Blesa, Zaragoza, Spain
- 6 Primary Health Care Center “Delicias Sur”, Zaragoza, Spain

Table 4 Frequency of signs and symptoms in symptomatic patients by age group

	< 1 y/o	1–4	5–9	10–14	Total
Fever	166/215 (77.2%)	431/665 (64.8%)	471/833 (56.5%)	587/1138 (51.6%)	1655/2851 (58.1%)
Cough	144/220 (65.5%)	364/685 (53.1%)	360/850 (42.4%)	499/1174 (42.5%)	1367/2929 (46.7%)
Nasal congestion	116/220 (52.7%)	310/685 (45.3%)	260/850 (30.6%)	369/1174 (31.4%)	1055/2929 (36.0%)
Odynophagia	26/218 (11.9%)	86/679 (12.7%)	218/850 (25.7%)	327/1174 (27.9%)	657/2921 (22.5%)
Headache	5/216 (2.3%)	32/680 (4.7%)	224/850 (26.4%)	418/1174 (35.6%)	679/2920 (23.3%)
Dyspnea	7/220 (3.2%)	21/685 (3.1%)	38/850 (4.5%)	67/1174 (5.7%)	133/2929 (4.5%)
Nauseas/vomits	20/220 (9.1%)	59/685 (8.6%)	60/850 (7.1%)	56/1174 (4.8%)	195/2929 (6.7%)
Diarrhea	40/220 (18.2%)	117/685 (17.1%)	93/850 (10.9%)	123/1174 (10.5%)	373/2929 (12.7%)
Abdominal pain	18/217 (8.3%)	67/680 (9.9%)	125/850 (14.7%)	108/1174 (9.2%)	318/2921 (10.9%)
Skin manifestations	3/220 (1.4%)	36/685 (5.3%)	31/850 (3.6%)	26/1174 (2.2%)	96/2929 (3.3%)
Malaise/irritability	21/217 (9.7%)	60/681 (8.8%)	104/850 (12.2%)	230/1174 (19.6%)	415/2922 (14.2%)
Myalgia	0/217 (0.0%)	15/679 (2.2%)	77/850 (9.1%)	146/1174 (12.4%)	238/2920 (8.2%)
Anosmia/dysgeusia	1/216 (0.5%)	8/678 (1.2%)	46/850 (5.4%)	149/1175 (12.7%)	204/2920 (7.0%)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.